



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Whence, $\phi = (\pi/2) - (\theta/2)$. Also

$$R = \frac{[1 + (dy/dx)^2]^{-\frac{3}{2}}}{d^2y/dx^2} = 4a \sin \theta/2$$

Hence, $P = v_0^2 \div (4a \sin \theta/2) - g \sin \theta/2$. But, since $y = \theta(1 - \cos \theta)$,

$$2 \sin^2 \frac{\theta}{2} = \frac{y}{a}.$$

Hence,

$$P = \frac{v_0^2 - 4ag \sin^2 \theta/2}{4a \sin \theta/2} = \frac{v_0^2 - 2gy}{4a \sin \theta/2},$$

But, at the point where the hound tumbled through, $y = y_0 = (v_0/2g)^2$. Hence, at that point, $P = 0$.

NUMBER THEORY.

212. Proposed by C. N. SCHMALL, New York City.

Given any positive integer N greater than 1; to prove that the sum of all the positive integers less than N and prime to N equals $\frac{1}{2}N \cdot \phi(N)$.

SOLUTION BY H. C. FEEMSTER, York College, Nebraska.

Let $N = a^h b^k c^l \dots$, where a, b, c, \dots are primes. Then

$$\phi(N) = N \cdot \frac{a-1}{a} \cdot \frac{b-1}{b} \cdot \frac{c-1}{c} \cdot \dots = N \left[1 - \Sigma \frac{1}{a} + \Sigma \frac{1}{ab} - \Sigma \frac{1}{abc} + \dots \right],$$

the number of positive integers less than N and prime to N .

For every number $p < N/2$ and prime to N , there is a number $N - p > N/2$ and less than N and prime to N . Now there are $\phi(N)$ of these numbers or $\frac{1}{2}\phi(N)$ pairs of these numbers. But the sum of each pair is N , so the entire sum is $\frac{1}{2}N \cdot \phi(N)$, as required.

213. Proposed by R. D. CARMICHAEL, Indiana University.

Prove that no relatively prime integers x and y exist such that the difference of their fourth powers is a perfect cube.

SOLUTION BY ELIJAH SWIFT, University of Vermont.

We are to prove the non-existence of any equation $x^4 - y^4 = z^3$. There are two cases to consider: (1) x and y both odd; (2) one even and one odd.

In the second case $x^2 + y^2$ and $x^2 - y^2$ are prime to each other, since any common factor would divide their sum $2x^2$, and difference $2y^2$; but these have, by hypothesis, 2 as their only common factor, and 2 is not a factor of $x^2 + y^2$ because x is even and y odd, or vice versa.

Since the product of $x^2 + y^2$ and $x^2 - y^2$ is a perfect cube, each of the factors must be, say, $x^2 + y^2 = a^3$, $x^2 - y^2 = b^3$. Call $2xy = c$. Then $a^6 - b^6 = c^2$, which is impossible. (See Number Theory, problem 209 in the March, 1914, MONTHLY which denies the existence of such an equation.)